

$\chi_{c0}(3860)$ $I^G(J^{PC}) = 0^+(0^{++})$

OMMITTED FROM SUMMARY TABLE

The assignment $J^P = 0^+$ is preferred over 2^+ by 2.5 sigma.Observed by CHILIKIN 17 using full amplitude analysis of the process
 $e^+ e^- \rightarrow J/\psi D\bar{D}$, where $D = D^0, D^+$. **$\chi_{c0}(3860)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3862 $+26$ -32	CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D\bar{D}$

 $\chi_{c0}(3860)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
201 $+154$ -67	CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D\bar{D}$

 $\chi_{c0}(3860)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^0 \bar{D}^0$	seen
Γ_2 $D^+ D^-$	seen

 $\chi_{c0}(3860)$ BRANCHING RATIOS

$\Gamma(D^0 \bar{D}^0)/\Gamma_{\text{total}}$	Γ_1/Γ
seen	CHILIKIN 17 BELL $e^+ e^- \rightarrow J/\psi D^0 \bar{D}^0$

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$	Γ_2/Γ
seen	CHILIKIN 17 BELL $e^+ e^- \rightarrow J/\psi D^+ D^-$

 $\chi_{c0}(3860)$ REFERENCESCHILIKIN 17 PR D95 112003 K. Chilikin *et al.* (BELLE Collab.) JPC